


Managing your insulin doses during illness if you have type 1 or type 2 diabetes¹

Diabetes:

What to do when you are ill



- How illness affects your blood glucose levels
- Monitoring your diabetes
- Managing your medication
- What to eat and drink
- When to seek help

The content was generated independently by TREND-UK in conjunction with SB Communications Group. AstraZeneca funded the printing.

How does illness affect your blood glucose?

- When you are ill, or have an infection or a virus such as a cold, your blood glucose levels may rise in response to this, even if you are not eating
- This may make you feel thirsty and pass urine more frequently, which can make you dehydrated. You may therefore need to increase the dose of your diabetes medication to combat this
- If you have type 1 diabetes, your body can produce a substance called ketones when you are unwell, which can cause a serious condition called diabetic ketoacidosis
- It is possible to manage your diabetes during illness effectively and keep your blood glucose levels down by following the simple advice given in this leaflet

DO NOT STOP TAKING YOUR INSULIN

General guidance for people with diabetes

- Try to drink at least 4 to 6 pints (2.5 to 3.5 L) of sugar-free fluid in 24 hours to prevent becoming dehydrated
- Avoid strenuous exercise
- Treat symptoms such as a high temperature or a cough with basic medicines such as painkillers and cough medicines. These do not have to be sugar-free varieties, as they are taken in small quantities
- See your GP if you think you have an infection, as you may need antibiotics
- Test your blood or urine glucose levels, checking them at least four times a day¹
- If you are on insulin and have been taught how to adjust, increase or decrease your doses accordingly, see the flowchart on the back pages¹
- Seek medical help if your readings remain higher than usual, you feel unwell and you are not sure what to do

Meal replacement suggestions

Try to eat as normal but if you cannot manage your usual meals, replace these with light and easily digested foods such as soups and milky puddings. See the table below for a list of alternative food options.

Type of food alternative	Amount (Each serving provides approximately 10 g of carbohydrate)		
Lucozade™ Energy	50 mL	2 fl oz	¼ glass
Fruit juice*	100 mL	4 fl oz	½ glass
Cola (NOT diet)*	100 mL	4 fl oz	½ glass
Lemonade (NOT diet)*	150–200 mL	5–7 fl oz	¾–1 glass
Milk	200 mL	7 fl oz	1 glass
Soup*	200 mL	7 fl oz	1 mug
Ice cream*	50 g	2 oz	1 large scoop
Complan®	–	–	3 level teaspoons (as a drink)
Drinking chocolate*	–	–	2 level teaspoons (as a drink)
Ovaltine® or Horlicks®	–	–	2 level teaspoons (as a drink)

*Sugar quantities may vary according to brand
Keep sugary drinks at home for emergencies

Call your GP, diabetes specialist nurse or healthcare professional for immediate help:

- If you are pregnant
- If you have a child or are the carer of a child with diabetes
- If you have type 1 diabetes and cannot keep fluids down
- If you have type 1 diabetes, have positive ketones and do not know how to adjust your insulin to manage these
- If, despite taking the advice in this leaflet, your symptoms are getting worse

Managing your diabetes medication If you have type 2 diabetes

If you have type 2 diabetes and are taking diabetes medication:

- Continue to take your tablets even if you are not eating
- If you are taking metformin or an SGLT2 inhibitor and you are vomiting or have diarrhoea, you should stop this medication
- You may need to increase the dose of your tablets or even need insulin injections for a short time while you are ill²
- Contact your GP if you are able to check your blood glucose levels and find they are constantly higher than 15 mmol/L
- If you develop abdominal pain, nausea and vomiting, stop the medication **immediately** and seek medical attention

If you have type 2 diabetes and take insulin as well as diabetes medication:

- Monitor and record your blood glucose levels at least four times a day (at mealtimes even if you are not eating, and at bedtime)
- If your blood glucose levels are **higher** than usual, you may need to increase your insulin (see overleaf). Contact your GP or Diabetes Nurse if you are not sure how to do this¹
- If your blood glucose levels are **lower** than usual (because you are eating less than usual) you may need to reduce your insulin dose¹

Further information

- Diabetes UK Careline: 0845 120 2960
- Diabetes UK: www.diabetes.org.uk

References

1. TREND-UK (2013) *Managing diabetes during intercurrent illness in the community*. Available at: <http://trend-uk.org> (accessed 23.05.2013)
2. Diabetes UK (2013) *When you are ill*. Available at: http://www.diabetes.org.uk/Guide-to-diabetes/Living_with_diabetes/illness/When_you_are_ill/ (accessed 06.06.2013)